ELSEVIER

Palaeogeography, Palaeoclimatology, Palaeoecology 154 (1999) 383–384

PALAEO

Contents Volume 154, 1999

Special Issue: Biotic Recoveries from Mass Extinctions
edited by D.H. Erwin

Foreword 1

The generative phase and the first radiation event in the Early Silurian monograptid history
T. Koren' and M. Bjerreskov 3

Famennian and Tournaisian recoveries of shallow water Rugosa following late Frasnian and late Strunian major crises, southern Belgium and surrounding areas, Hunan (South China) and the Omolon region (NE Siberia)
E. Poty 11

Palaeoenvironments and faunal recovery after the end-Permian mass extinction
R.J. Twitchett 27

The early Toarcian (Early Jurassic) and the Cenomanian–Turonian (Late Cretaceous) mass extinctions: similarities and contrasts
P.J. Harries and C.T.S. Little 39

Benthic extinction and recovery patterns at the K/T boundary in shallow water carbonates, Denmark
E. Håkansson and E. Thomsen 67

Lower Danian bivalves, Stevns Klint, Denmark: continuity across the K/T boundary
C. Heinberg 87

Silica-secreting biota and mass extinctions: survival patterns and processes
G. Racki 107

Research Papers

Milankovitch-scale palaeoclimate changes in pale–dark bedding rhythms from the Early Cretaceous (Hauterivian and Barremian) of eastern England and northern Germany
J. Mutterlose and A. Ruffell 133

Palaeomagnetic chronology of the evaporitic sedimentation in the Neogene Fortuna Basin (SE Spain): early restriction preceding the 'Messinian Salinity Crisis'
J. Dinarès-Turell, F. Ortí, E. Playà and L. Rosell 161

Three large shifts in East Asian monsoon circulation indicated by loess–paleosol sequences in China and late Cenozoic deposits in Japan
J. Xiao and Z. An 179

Pacific Ocean intermediate and deep water circulation during the Pliocene
P.B. Kwiek and A.C. Ravelo 191

Palaeoclimatic and chronostratigraphic interpretations from strontium, carbon and oxygen isotopic ratios in molluscan fossils of Quaternary Seoguipo and Shinyangri Formations, Cheju Island, Korea
K.H. Kim, T. Tanaka, T. Nakamura, K. Nagao, J.S. Youn, K.R. Kim and M.Y. Yun 219

Astronomical calibration of loess–paleosol deposits at Luochuan, central Chinese Loess Plateau
H. Lu, X. Liu, F. Zhang, Z. An and J. Dodson 237

Paleoenvironmental changes across the Cretaceous–Tertiary boundary at Koshak, Kazakhstan, based on planktic foraminifera and clay mineralogy
A. Pardo, T. Adatte, G. Keller and H. Oberhänsli 247
Anoxia pre-dates Frasnian–Famennian boundary mass extinction horizon in the Great Basin, USA
J.F. Bratton, W.B.N. Berry and J.R. Morrow 275
Calcium carbonate dissolution rates in deep-sea bivalve shells on the East Pacific Rise at 21°N: results of an 8-year in-situ experiment
M.J. Kennish and R.A. Lutz 293
Faunal remains from a Middle Pleistocene lacustrine marl in Dakhleh Oasis, Egypt: palaeoenvironmental reconstructions
C.S. Churcher, M.R. Kleindienst and H.P. Schwarcz 301
Temporal change of life-history traits in fossil bivalves: an example of *Phacosoma japonicum* from the Pleistocene of Japan
S. Sato 313
The shell of the Queen Scallop *Aequipecten opercularis* (L.) as a promising tool for palaeoenvironmental reconstruction: evidence and reasons for equilibrium stable-isotope incorporation
J.A. Hickson, A.L.A. Johnson, T.H.E. Heaton and P.S. Balson 325
Environmental history of the German Lower Rhine Embayment during the Middle Miocene as reflected by carbon isotopes in brown coal
A. Lücke, G. Helle, G.H. Schleser, I. Figueiral, V. Mosbrugger, T.P. Jones and N.P. Rowe 339

1999 Subject Index, Volumes 144–154 353

Author Index Volume 154 381